

The Effectiveness of Body Percussion Rhythmic Exercises on Motor Skills in Children with Mild Intellectual Disability Between 8-12 Years Old

Salehe lotfi¹, Mitra Khalafbeigi^{1*}, Nader Matin Sadr², Seyed Hassan Saneii³

1. Department of Occupational Therapy, School of Rehabilitation Sciences, Iran University of Medical Sciences, Tehran, Iran
2. Department of Occupational Therapy, Mashhad University of Medical Sciences, Mashhad, Iran
3. Department of Basic Sciences in Rehabilitation, School of Rehabilitation Sciences, Iran University of Medical Sciences, Tehran, Iran

Article Info

Received: 2018/04/26

Accepted: 2018/08/18

Published Online: 2018/09/27

DOI: 10.30699/fdisj.1.3.1.40

How to Cite This Article

lotfi S, Khalafbeigi M, Matin Sadr N, Saneii S H. The Effectiveness of Body Percussion Rhythmic Exercises on Motor Skills in Children with Mild Intellectual Disability Between 8-12 Years Old. *Func Disabil J*. 2018; 1 (3) :40-47

Use your device to scan and read the article online



ABSTRACT

Background and Objective: Intellectual disability is a common neurological disorder that involves concomitant impairments in mental and adaptive functioning and begins during the developmental period. Percussion is a rhythmic movement activity and the art of body-shaping to produce a variety of sounds for educational, therapeutic, ethnological and social purposes. The purpose of this study was to investigate the effectiveness of percussion rhythmic exercises on motor skills of mild mentally disabled children.

Methods: This is a randomized clinical trial in which 60 children with mild intellectual disability aged 8 to 12 years were selected by convenience sampling and divided into two groups: experimental (n = 31) and control (n = 29). The experimental group received percussion inflatable exercises as a group for 12 weeks, 2 sessions per week and 30 minutes per session and the control group received only the usual school schedule. Both groups were evaluated before and after by Bruininks-oseretsky Test of Motor Proficiency. Paired T-test was used for intra-group comparisons and independent T-test was used to compare the two groups.

Results: Results indicated that Body Percussion Rhythmic Exercises had a significant effect on some motor items including: fine motor skills, bilateral coordination, upper extremity coordination, motor visual control, speed and upper extremity skills.

Conclusion: Therefore, it can be stated that group Body Percussion Rhythmic Exercises is a good way to improve motor skills in children with mild intellectual disability.

Keywords: Rhythm, Body Percussion, Motor skills, Mild Intellectual disability

Corresponding information:

Mitra Khalafbeigi, Department of Occupational Therapy, School of Rehabilitation Sciences, Iran University of Medical Sciences, Tehran, Iran. Email:khalafbeigi.m@iums.ac.ir Tel: +98-21-

Copyright © 2018, Function and Disability Journal. This is an open-access article distributed under the terms of the Creative Commons Attribution-noncommercial 4.0 International License which permits copy and redistribute the material just in noncommercial usages, provided the original work is properly cited.

Introduction

According to the DSM-5, intellectual disability is a disorder that begins during the developmental period and involves simultaneous deficits in mental functioning and consistent functioning in conceptual, social, and practical contexts (Association, 2013). In Iran,

the intellectually disabled constitute 1% to 3% of the population (Nesayan & Gandomani, 2016) with a mild or educable type accounting for more than 66% of the population (Ghamari, Rafeei, Soltani, & Ghamari, 2016). These individuals have deficits in several areas such as communication, daily living skills, and motor

41. Effectiveness of Body Percussion Rhythmic Exercises on Motor Skills

Table 1. Bruininks-oseretsky Test of Motor Proficiency

R	L	R	L
★	★	★	★
✚	✚	✚	✚
★	★	★	★

skills (Daftari, Behnia, Rassafiani, Sajedi, & Biglarian, 2014). Motor skills play a very important role in learning and provide the basis for the development of other important learning such as educational and social skills (Pahlevanian, Rasoolzadeh, & Khalily, 2012). To correct these, occupational therapists use a variety of interventions, one of which is music activity.

One of the subsections of music education is “music and movement”. Body Percussion is a fringe discussion of the different types of rhythm exercises in which the body presents itself as an audio element (Ahokas, Burger, & Thompson, 2014) and as the first musical instrument from which we can obtain an infinite range of training resources through verbal beats and sounds that create organized skills and mastery of movement (FJ Romero-Naranjo, 2013). This method has been chosen as a training method and can be run without any additional tools and can be easily ported to any accessible environment (Ahokas et al., 2014). Also, it can be performed individually, in paired groups or in groups with more than 3 people (in a circle, a semicircle, square or linear). Each of the aforementioned layouts stimulates individuals’ physical and social interaction in a different way (F. J. Romero-Naranjo, 2014). Various limb movements, such as clapping, slapping, kicking, breaking and stomping, produce sounds with varying amplitudes that generate rhythm. Percussion training enhances the nervous system’s sensitivity, rhythm knowledge about time and space and its expression (Ahokas et al., 2014). Importantly, different cortical and subcortical regions in both hemispheres are stimulated during these activities.

As such, movement affects the prefrontal cortex, cerebellum, and basal nuclei. To properly perform body movements, cognitive aspects such as memory and attention are involved, which itself activates various areas of the brain, including the frontal cortex, cerebellum, and hippocampus. The motivational aspects of

learning new skills also stimulate the limbic system, which in turn enhances learning (Ghaffarian, 2014). Group body percussion exercises facilitate direct experience-based learning (learning by doing) and provide an opportunity and space for individuals to creatively and collectively learn group skills (A. Romero-Naranjo, Romero Naranjo, & Bofill, 2016). In addition to developing such skills in the school environment, they contribute to the overall educational, individual and social development and provide opportunities for increased creativity and exploratory behavior, friendships that facilitate individual maturity and abilities related to learning and motivation (Moral-Bofill, Romero-Naranjo, Albiar-Aliaga, & Cid-Lamas, 2015). Research show that applying this method improves cognitive skills, spatial vision, psychomotor (including awareness of one’s own body, hand-eye coordination, writing skills, lateral superiority) (A. Carretero-Martínez & Romero-Naranjo, 2015), executive functions (Cozzutti, Guaran, Blessano, & RomeroNaranjo, 2017), socioemotional stimuli (Marcuzzi & Romero-Naranjo, 2017), and friendly relationships (Fabra-Brell & Romero-Naranjo, 2017). The present study investigated the effectiveness of this method in children with mild intellectual disability.

Materials and Methods

Trial Design

This study is a randomized clinical trial, with code of ethics IR.IUMS.REC.1397.1132, and trial ID 39719. Bruininks-oseretsky Test of Motor Proficiency was performed by a blinded assessor before and after the intervention.

Randomization

Initially two schools that were willing to cooperate with the research project were selected through available sampling and sixty students with inclusion criteria were enrolled in the study and then randomly assigned to two groups of experimental (n=31) and control

Table 2. Comparison of mean and standard deviation of variables with normal distribution before and after treatment using paired T-test

Variables	Group	Number	Mean	Standard Deviation	T	Significance Level	
Balance	Test	Pre-test	29	23.2414	7.69348	-4.429	0.000
		Post-test		27.4483	5.57152		
	Control	Pre-test	28	17.1786	10.24895		
		Post-test		19.0714	10.44537		
Two-way coordination	Test	Pre-test	29	6.6207	2.45552	-7.616	0.000
		Post-test		9.6207	2.59689		
	Control	Pre-test	28	5.2143	3.22441		
		Post-test		5.7500	3.63751		
Power	Test	Pre-test	29	9.4828	4.05868	-8.885	0.000
		Post-test		13.1034	4.16915		
	Control	Pre-test	28	8.2143	5.10887		
		Post-test		9.5714	6.23270		
Upper limb coordination	Test	Pre-test	29	10.6207	4.67832	-5.547	0.000
		Post-test		13.8621	5.04072		
	Control	Pre-test	28	9	5.46368		
		Post-test		9.5357	6.15529		
Motor vision control	Test	Pre-test	29	13	4.40779	-7.045	0.000
		Post-test		16.6552	2.66292		
	Control	Pre-test	28	10.5000	4.02308		
		Post-test		12.1429	4.24015		
Upper extremity speed and skill	Test	Pre-test	29	16.1034	5.62739	-8.033	0.000
		Post-test		21.8621	5.65511		
	Control	Pre-test	28	13.7143	5.58342		
		Post-test		16.6429	6.09927		
Coarse motor skills	Test	Pre-test	29	43.7586	14.15641	-10.698	0.000
		Post-test		57.2069	11.80913		
	Control	Pre-test	28	33.8571	19.65200		
		Post-test		38.7857	21.10881		
Fine motor skills	Test	Pre-test	29	30.2069	9.75185	-9.979	0.000
		Post-test		40.1034	7.71522		
	Control	Pre-test	28	25.2500	9.7282		
		Post-test		30	10.59000		

43. Effectiveness of Body Percussion Rhythmic Exercises on Motor Skills

Table 3. Intra-group comparison of variables with abnormal distribution before and after treatment using Wilcoxon test

Variables	Groups	Number	Z	Significance Level
Sprint and agility	Test	29	-1.802	0.072
	Control	28		
Response speed	Test	29	-0.963	0.335
	Control	28		

Table 4. Comparison of mean and standard deviation of variables with normal distribution in two groups of test and control using independent T-test

Variables	Groups	Number	Z	Significance Level	
Sprint and agility	Test	Pre-test	29	-3.930	0.000
		Post-test			
	Control	Pre-test	28	-2.740	
		Post-test			
Response speed	Test	Pre-test	29	-2.122	0.034
		Post-test			
	Control	Pre-test	28	-1.508	
		Post-test			

Table 5. Comparison of variables with abnormal distribution in two groups of test and control using Mann-Whitney U test

Variables	Groups	Number	Mean	Standard Deviation	T	Significance Level	
Balance	/Pre-test	Test	29	4.2069	5.11566	1.970	0.054
	Post-test	Control	28	1.8929			
Two-way coordination	/Pre-test	Test	29	3	2.12132	5.100	0.000
	Post-test	Control	28	0.5357			
Power	/Pre-test	Test	29	3.7931	2.25799	3.878	0.000
	Post-test	Control	28	1.3571			
Upper limb coordination	/Pre-test	Test	29	3.2411	3.14705	3.289	0.002
	Post-test	Control	28	0.5357			
Motor vision control	/Pre-test	Test	29	3.6552	2.79382	2.886	0.006
	Post-test	Control	28	1.6429			
Upper extremity speed and skill	/Pre-test	Test	29	5.7586	3.86056	2.744	0.008
	Post-test	Control	28	2.9286			
Coarse motor skills	/Pre-test	Test	29	13.5862	6.54390	5.576	0.000
	Post-test	Control	28	4.9286			
Fine motor skills	/Pre-test	Test	29	9.8968	5.34085	3.850	0.000
	Post-test	Control	28	4.7500			

(n=29). The mean age of the test group was 9.6 and the mean age of the control group was 9.4.

Participants

In the schools in question, inclusion criteria were considered while selecting 60 samples. Inclusion criteria for all participants included: Diagnosis of mild intellectual disability (educable) according to the Department of Special Education, age range of 8-12 years, no neurological disorders such as cerebral palsy, epilepsy and seizures, lack of associated psychiatric disorders such as attention deficit / hyperactivity disorder and autism, lack of orthopedic disorders that cause problems in child movement, and lack of visual and hearing impairment that impair understanding of instructions. After selecting the samples, the project was described to the targets' parents and teachers and demographic questionnaire and consent form were completed. Exclusion criterium was no lack of cooperation of the students or their family. By the end of the intervention, after omitting 2 cases the experimental group had 29 cases and the control group had 28 cases after omitting only 1 sample.

Setting

This study was conducted in Kashefi Kashani Mashhad boys and girls schools in autumn and winter of 2018.

Intervention

Both groups had traditional school education as the underlying treatment, and the experimental group received music percussion interventions in groups for 12 weeks and two 30 minutes sessions per week. Students would visit between 8 AM and 12 AM during the designated groupings for practice. The exercises were researcher-made and selected from simple to complex, based on team experience, resource review, and under the supervision of rhythmic instrument experts. The focus was on teamwork and included aspects of social skills such as teamwork, ability to adapt to the group, play and leisure, eye contact, expressing emotion, and actively participating in the group. Four percussions were selected which included: clapping, hitting on the thigh, Hitting on the chest, stamping the feet. A symbol was defined for each percussion and arranged in

tables to practice as a template. For example, the thigh was defined with the star symbol and the chest with the cross symbol. In this case, the exercise can be done in Table 1.

It is one of the most commonly used tests in motor performance assessment and is a standard reference test that measures motor skills of 4.5 to 14.5 years old children and consists of eight sub-tests with 46 items. It takes 45-60 minutes to complete the whole test. The full test consists of eight sub-tests including four sub-tests, coarse motor skills (sprinting, balance, two-way coordination, power), one upper-body coordination skills test, and three skills test sub-tests. They evaluate fine motor (response speed, motor vision control, upper extremity speed and skill). The validity of this test was 0.78, the inter-rater reliability was 0.99 - 0.92 before and after the test, 0.80 to 0.80 (Soltanikhadiv, Kamali, Rafiei, & Taghizade, 2014).

Statistical Method

Data were analyzed by SPSS 21 (SPSS Inc., Chicago, IL. USA) (significance level $P < 0.05$). Normality of data distribution was assessed by Kolmogorov-Smirnov test. Paired t-test (Table 2) and Wilcoxon (Table 3) were used to compare within-group scores. Independent T-test (Table 4) and Mann-Whitney U test (Table 5) were used to compare the scores of the two groups.

Results

According to the results (Tables 2-5), there is a significant difference in the variables of bilateral coordination, upper extremity coordination, visual-motor control, upper extremity speed and skill, coarse and fine motor skills, meaning that intervention of body percussion rhythmic exercises were effective on these motor items. But in the items of balance, speed and agility and speed of response, there was no significant difference between groups.

Discussion

The findings of the study confirmed that a percussion rhythmic program based on the group participation of students in school has a significant effect on some motor items. Various studies have investigated the efficacy of body percussion on eye-hand coordination (A. Car-

45. Effectiveness of Body Percussion Rhythmic Exercises on Motor Skills

retero-Martínez & Romero-Naranjo, 2015; Colomino & Naranjo, 2014), organ coordination (Cozzutti, Blessano, Biaggio, Tomasin, & Romero-Naranjo, 2017; Ahokas et al., 2014) and confirmed the improvement of psychomotor skills (Andrea Carretero-Martínez, Romero-Naranjo, PonsTerrés, & Crespo-Colomino, 2014). Body percussion activities play an important role in improving children's learning by activating all areas of the brain including cognitive processing, attention types, muscular and rhythmic memory, fine motor skills, and emotional and social aspects (Colomino & Naranjo, 2014). This is in line with the results of this study. Results also showed that percussion rhythmic exercises did not have a significant effect on sprinting items, balance and response speed. To increase speed, one must practice the ability to use maximum force during very fast movements.

In the present study, the focus of the exercises was on proper percussion on the target organs and producing rhythmic sound from the body. The same is true of agility and balance. Since these children are slower than their peers in initiating and performing target movements, reaction times, and movement times (Kahrizsangi, Salehi, & Heydari, 2012) and need to spend longer time on learning, accelerating and dealing with agility and balance was virtually impossible. In addition, by searching for various articles in the field of percussion or rhythmic movements, no study was found to confirm the effectiveness of such exercises on sprinting and agility. Inconsistent studies of

References

Ahokas, R., Burger, B., & Thompson, M. (2014). Brain and Body Percussion: The relationship between motor and cognitive functions Paper presented at the Proceedings of the 7th International Conference of Students of Systematic Musicology (SysMus14), London.

Association, A. P. (2013). Diagnostic and statistical manual of mental disorders : DSM-5 (Fifth ed.). Arlington.

Carretero-Martínez, A., & Romero-Naranjo, F. J. (2015). Stimulation of Fine Psychomotor Skills in Children. Methodological Introduction According to the BAPNE Method International Journal of Innovation and Research in Educational Sciences, 2(6), 2349–5219

response speed, or speed of action, include the 2017 Cozzotti study showing that percussion stimulation in training, improves motor sequence retention and more tasks simultaneously (Cozzutti, Guaran, et al., 2017) Also, a 2014 study by Colomino and Naranjo confirms that body percussion rhythmic exercises can help the process of recovery of dyslexic individuals in terms of speed of action (Colomino & Naranjo, 2014). To justify the insignificance of items for sprinting, balance, and response speed, the 24-session intervention period seems to be too short to improve for educated children or the time allocated to each session (30 to 45 minutes) is insufficient.

Conclusion

The body percussion method is one of the types of stimuli that is physically non-terminable and almost readily available. Therefore, it can be used as an effective and enjoyable therapeutic modality in a variety of sectors, including rehabilitation of intellectually disabled children and motor retardation, in accordance with scientific principles.

Acknowledgements

The authors would like to thank the families of all the children who participated in this study. Also they appreciate the help of the staff of the schools where this study was conducted, for their cooperation.

Conflict of Interest

No conflict of interest was declared by the authors.

Carretero-Martínez, A., Romero-Naranjo, F. J., PonsTerrés, J. M., & Crespo-Colomino, N. (2014). Cognitive, visual-spatial and psychomotor development in students of primary education through the body percussion –BAPNE Method Procedia - Social and Behavioral Sciences, 152, 1282 – 1287 doi:10.1016/j.sbspro.2014.09.363

Colomino, N. C., & Naranjo, F. J. R. (2014). Body percussion and dyslexia. Theoretical and practical contribution through the BAPNE method Procedia - Social and Behavioral Sciences, 132, 686 – 690 doi:10.1016/j.sbspro.2014.04.373

Cozzutti, G., Blessano, E., Biaggio, C. D., Tomasin,

- B., & Romero-Naranjo, F. J. (2017). Body solfege in the BAPNE method – Measures and divisions *Procedia - Social and Behavioral Sciences*, 237, 1572 – 1575 doi:10.1016/j.sbspro.2017.02.247
- Cozzutti, G., Guaran, F., Blessano, E., & RomeroNaranjo, F. J. (2017). Effects on executive functions in the BAPNE Method; a study on 8-9 years old children in Friuli Venezia Giulia, Italy *Procedia - Social and Behavioral Sciences* 237, 900 – 907 doi:10.1016/j.sbspro.2017.02.126
- Daftari, T., Behnia, F., Rassafiani, M., Sajedi, F., & Biglarian, A. (2014). Motor Deficits of Girls with Down Syndrome Comparing with Girls with Intellectual Disability in the School Ages. Special Issue “Pediatric Neurorehabilitation”, 14(6), 93-98.
- Fabra-Brell, E., & Romero-Naranjo, F. J. (2017). Body percussion: social competence between equals using the method BAPNE in Secondary Education (Design Research) *Procedia - Social and Behavioral Sciences*, 237, 1138 – 1142 doi:10.1016/j.sbspro.2017.02.168
- Ghaffarian, S. (2014). General anatomy with clinical notes. Mashhad: Mashhad University of medical sciences.
- Ghamari, N., Rafeei, S., Soltani, R., & Ghamari, Z. (2016). Effect of balance exercises along with gross movement on gross movements and balance function of children with mental retardation *Journal of Modern Rehabilitation*, 9(5), 69-76.
- Kahrizsangi, g. G., Salehi, h., & Heydari, I. (2012). The Effect of a Rhythmic Motor Program on Perceptual–Motor Abilities of Educable Mentally Retarded Children *A Journal of Faculty of Physical Education and Sport Sciences*, 4(1), 75-92. doi:10.22059/jmlm.2012.25076
- Marcuzzi, C., & Romero-Naranjo, F. J. (2017). BAPNE Method, Developmental Dyslexia and Inclusive Education: Cognitive, Socio-Emotional and Psychomotor Stimulation in Secondary School. A Practical Resource for Education within a Cross Curriculum. *Procedia - Social and Behavioral Sciences* 237, 1291 – 1298 doi:10.1016/j.sbspro.2017.02.212
- Moral-Bofill, L., Romero-Naranjo, F. J., Albiar-Aliaga, E., & Cid-Lamas, J. A. (2015). The BAPNE Method as a School Intervention and Support Strategy to Improve the School Environment and Contribute to Socioemotional Learning (SEL). *International Journal of Innovation and Research in Educational Sciences*, 2(6), 2349–5219
- Nesayan, A., & Gandomani, R. A. (2016). Effectiveness of Social Skills Training on Behavioral Problems in Adolescents with Intellectual Disability *Journal of Rehabilitation*, 17(2), 158-167.
- Pahlevanian, A. A., Rasoolzadeh, M., & Khalily, M. A. (2012). Comparison between normal and mental retard children with mental aged 6-7 on motor skills. *Koomesh*, 13(4), 460 - 464.
- Romero-Naranjo, A., Romero Naranjo, F. J., & Bofill, L. (2016). Body Percussion and Team Building through the BAPNE Method. *SHS Web of Conferences*, 26, 01020. doi:10.1051/shsconf/20162601020
- Romero-Naranjo, F. J. (2013). Science & art of body percussion: a review *JOURNAL OF HUMAN SPORT & EXERCISE*, 8(2), 442-457.
- Romero-Naranjo, F. J. (2014). Body percussion and memory for elderly people through the BAPNE method *Procedia - Social and Behavioral Sciences* 132, 533 – 537 doi:10.1016/j.sbspro.2014.04.349
- Soltanikhadiv, K., Kamali, M., Rafiei, S., & Taghizade, G. (2014). Relationship between the Bruininks Oseretsky Test of Motor Proficiency and the Peabody Developmental Motor Scale on Children with EducableIntellectual and Developmental Disabilities. *Journal of Research in Rehabilitation Sciences*, 10(2), 306-314.

بررسی اثربخشی تمرینات ریتمیک بادی پرکاشن بر مهارت‌های حرکتی کودکان کم‌توانی ذهنی خفیف ۸-۱۲ سال

صالحه لطفی^۱، میترا خلف بیگی^{۱*}، نادر متین صدر^۲، سید حسن صانعی^۳

۱. گروه آموزشی کاردرمانی، دانشکده علوم توانبخشی، دانشگاه علوم پزشکی ایران، تهران، ایران
۲. گروه آموزشی کاردرمانی، دانشکده علوم توانبخشی، دانشگاه علوم پزشکی مشهد، مشهد، ایران
۳. گروه آموزشی علوم پایه در توانبخشی، دانشکده علوم توانبخشی، دانشگاه علوم پزشکی ایران، تهران، ایران

چکیده	اطلاعات مقاله
<p>زمینه و هدف: کم‌توانی ذهنی یک اختلال شایع نورولوژیکی است که شامل نقص‌های هم‌زمان در عملکردهای ذهنی و سازگارانه است و در طول دوره رشدی شروع می‌شود. بادی پرکاشن یک فعالیت حرکتی ریتمیک و هنر ضربه‌زدن به بدن برای تولید انواع صداهای مختلف برای اهداف آموزشی، درمانی، قوم‌شناسی و اجتماعی است. هدف از این مطالعه بررسی اثربخشی تمرینات ریتمیک بادی پرکاشن بر مهارت‌های حرکتی کودکان کم‌توان ذهنی خفیف است.</p> <p>روش کار: این مطالعه کارآزمایی بالینی تصادفی است که ۶۰ کودک کم‌توان ذهنی خفیف ۸ تا ۱۲ سال به صورت در دسترس انتخاب شدند و به روش تصادفی به دو گروه آزمون (۳۱ نفر) و کنترل (۲۹ نفر) تقسیم شدند. گروه آزمون تمرینات بادی پرکاشن را در قالب گروهی برای ۱۲ هفته، ۲ جلسه در هفته و هر جلسه ۳۰ دقیقه دریافت کرد و گروه کنترل فقط برنامه رایج مدرسه را دریافت کرد. هر دو گروه به‌طور قبل و بعد توسط آزمون تب‌حرکتی برونینیکس ازورسکی مورد ارزیابی قرار گرفتند. از آزمون T زوج برای مقایسه درون گروهی و از T مستقل برای مقایسه دو گروه استفاده شد.</p> <p>یافته‌ها: نتایج حاکی از آن بود که تمرینات ریتمیک بادی پرکاشن تأثیر قابل توجهی در برخی آیتم‌های حرکتی از جمله: مهارت‌های حرکتی درشت و ظریف، هماهنگی دو طرفه، هماهنگی اندام‌های فوقانی، کنترل بینایی حرکتی، سرعت و مهارت اندام فوقانی دارد.</p> <p>نتیجه‌گیری: بنابراین می‌توان اظهار کرد که اجرای گروهی تمرینات ریتمیک بادی پرکاشن یک راه مناسب برای بهبود مهارت‌های حرکتی در کودکان کم‌توان ذهنی خفیف است.</p> <p>واژه‌های کلیدی: ریتم، بادی پرکاشن، مهارت‌های حرکتی، کم‌توانی ذهنی خفیف</p>	<p>تاریخ وصول: ۱۳۹۷/۰۲/۰۶</p> <p>تاریخ پذیرش: ۱۳۹۷/۰۵/۲۷</p> <p>انتشار آنلاین: ۱۳۹۷/۰۷/۰۵</p> <p>نویسنده مسئول:</p> <p>میترا خلف بیگی</p> <p>گروه آموزشی کاردرمانی، دانشکده علوم توانبخشی، دانشگاه علوم پزشکی ایران، تهران، ایران</p> <p>پست الکترونیک: khalafbeigi.m@iums.ac.ir</p> <p>تلفن: +۹۸-۲۱-</p>